

Complementary Tolls in the periodontium: how periodontal bacteria modify complement and Toll-like receptor responses to prevail in the host

JENNIFER L. KRAUSS, JAN POTEPA, JOHN D. LAMBRIS & GEORGE HAJISHENGALLIS

Innate immunity is a phylogenetically ancient system of host defense and represents the inherited resistance to infection (75). Until relatively recently, the innate immune response was viewed as a nonspecific and temporary expedient to 'buy time' until the activation of adaptive immunity, which comprises the system of B and T lymphocytes, each of which expresses antigen receptors of exquisite specificity (41). Although lacking the ability to make such fine structural distinctions, innate immunity is nevertheless endowed with considerable specificity. Indeed, germline-encoded receptors (collectively known as pattern-recognition receptors) can detect and respond to conserved and generally distinct microbial structures that are shared by related groups of microorganisms (e.g. lipopolysaccharide of gram-negative bacteria or lipoteichoic acid of gram-positive bacteria) (105). Most importantly, innate immunity is sophisticated enough to make judgments that instruct the initiation and progression of the adaptive immune response (41, 105). In this regard, the acquired specificity of the antigen receptors is not the result of co-evolution with microbes but the outcome of randomly generated gene recombination. Thus, even though the adaptive immune receptors can bind virtually any structure, they have no clue on the biological context of the encountered antigen (i.e. should they respond or not?). This information, however, is provided by innate immune mechanisms, which act as mediators between detection of infection and induction of the adaptive response. Not surprisingly, therefore, suc-

cessful pathogens, which disarm or subvert host defenses, target preferentially innate immunity (42) and particularly central systems such as the complement system and the Toll-like receptor (TLR) family of pattern-recognition receptors (96, 140).

In the oral cavity, innate immunity contributes significantly to antimicrobial defense, although inadequate or over-exuberant activation of the innate response may lead to oral disease, such as periodontitis (36, 53). In this context, periodontal health represents a dynamic state where proinflammatory and antimicrobial activities to control infection are optimally balanced by anti-inflammatory mechanisms to prevent unwarranted inflammation (43). This homeostatic balance may be disrupted, however, either by genetic defects in host immunity or by pathogens that undermine host-defense mechanisms (43, 89, 93). It should be noted that pathogen-instigated immune suppression of specific pathways and destructive inflammatory responses in the periodontium are not necessarily mutually exclusive, because the latter may arise as a consequence of the inability to control infection (52).

This review focuses on two key innate immune components – the complement system and the TLRs – and discusses how these systems interact with each other and with periodontal bacteria. The ability of periodontal pathogens to persist and establish chronic infections suggests that they may have evolved ways to disarm these defense mechanisms or subvert them to their advantage. Understanding the

mechanisms of periodontal host–pathogen interplay can offer important insights into the disease pathogenesis and facilitate the rational design of therapeutic interventions. A brief background on complement and TLR biology is presented below to facilitate the discussion on their interactions with periodontal bacteria.

Complement, TLRs and potential for crosstalk

The term ‘complement’ was coined by Paul Ehrlich in the late 1890s to describe a heat-sensitive activity in serum that is complementary to that of antibody in causing lysis of bacteria (153). In line with this early view, the complement has been traditionally considered as an antimicrobial enzyme system found in serum and inflammatory exudates such as the gingival crevicular fluid (7, 110, 123). However, it is now well appreciated that complement constitutes a fundamental component of innate immunity, by virtue of its ability to orchestrate critical events during immune and inflammatory responses, including the regulation of other innate or adaptive immune pathways (67, 98, 103, 183).

The triggering of the complement system involves the sequential activation and proteolytic cleavage of a series of serum proteins, leading to recruitment and activation of inflammatory cells, microbial opsonization and phagocytosis, and direct lysis of targeted pathogens (103). In addition to the serum components, the integrated complement system also includes membrane-bound regulators and receptors for interactions with various mediators of the immune system. Complement activation can proceed through three distinct mechanisms, namely the classical, lectin, or alternative pathways (103) (Fig. 1). All three pathways converge at a central step, involving activation of the third component of complement (C3) by pathway-specific C3 convertases (95, 103). The activation of the classical pathway is initiated by antigen–antibody complexes, whereas the lectin pathway is triggered through the interaction of a secreted pattern-recognition receptor (the mannose-binding lectin) with specific carbohydrate groups on the surface of a variety of microorganisms. To ensure a fast and immediate response to invading pathogens, the complement cascade is maintained at a low level of activity (‘tick-over’) by the so-called alternative pathway. This pathway is initiated by spontaneous hydrolysis of C3 to C3(H₂O), thereby inducing a conformational change that allows bind-




Fig. 1. Activation pathways of the complement system. All three pathways converge at a central step, involving activation of the third component of complement (C3) by pathway-specific C3 convertases. The classical pathway is initiated by antigen–antibody (Ag–Ab) complexes and requires the participation of C1, C2 and C4. The lectin pathway is triggered through interaction of the mannose-binding lectin (MBL) with specific carbohydrate groups on the surface of microorganisms. The alternative pathway is initiated by spontaneously hydrolyzed C3 [C3(H₂O)], which can thereby form a complex with factor B, followed by factor B cleavage by factor D and formation of the initial alternative pathway C3 convertase (103). Moreover, the alternative pathway can be induced by bacterial lipopolysaccharide and lipooligosaccharide in a properdin-dependent manner (86). Proteolytic cleavage of a series of proteins downstream of C3 leads to the generation of potent effector molecules. These include the anaphylatoxins C3a and C5a, which activate specific receptors (C3aR and C5aR, respectively), although C5a also interacts with the so-called C5a receptor-like 2 (C5L2), which is only modestly characterized (91). Additional effectors generated downstream of C3 are the opsonins C3b and iC3b, the latter of which coats microbes and promotes their phagocytosis by complement receptor-3 (CR3). In the terminal pathway, C5b initiates the assembly of the C5b–9 membrane attack complex (MAC), which in turn induces microbial cell lysis (103). Complement activation can also occur through crosstalk with other physiological pathways, such as the coagulation system, in which thrombin acts as a C5 convertase (71).

ing to complement factor B and formation of the initial alternative pathway C3 convertase. This results in the rapid propagation of the alternative pathway, as long as there is no sufficient negative regulation as normally occurs with nonself surfaces (e.g. bacteria). In addition to this mechanism, the alternative pathway can be induced by bacterial lipopolysaccharide and lipooligosaccharide molecules in a way that strictly requires the participation of the plasma protein properdin (86). The alternative pathway may




Fig. 2. Microbial ligand specificities of human Toll-like receptors (TLRs). Those TLRs (i.e. TLRs 1, 2, 4, 5 and 6) that recognize extracellular microbial structures are expressed on the host cell surface. TLR2, in cooperation with its signaling partners, TLR1 or TLR6, detect mostly microbial cell wall components, such as lipoproteins, lipoteichoic acid (LTA), fimbriae, or yeast zymosan (1, 12). TLR4 and TLR5 recognize lipopolysaccharide (LPS)

bacterial flagellin, respectively, whereas no ligand has been identified for TLR10. Those TLRs (i.e. TLRs 3, 7, 8 and 9) specializing in detecting viral or bacterial nucleic acids are expressed intracellularly on endocytic vesicles. TLR3 recognizes double-stranded viral RNA, TLR7 and TLR8 recognize single-stranded viral RNA and TLR9 detects microbial CpG DNA. LPS, lipopolysaccharide; LTA, lipoteichoic acid.

represent up to 80% of complement activation (103). In all three pathways, proteolytic cleavage of a series of proteins downstream of C3 leads to the generation of effector molecules, including opsonins (C3b, iC3b) and anaphylatoxins (C3a, C5a). The iC3b fragment is generated by further cleavage of microbe-attached C3b and mediates phagocytosis by complement receptor-3 (Fig. 1). The inflammatory anaphylatoxins C3a and C5a activate seven-transmembrane domain G-protein-coupled receptors, known as the C3a receptor and C5a receptor (CD88), respectively. A newly identified, but modestly characterized, alternative receptor for C5a is the so-called C5a receptor-like 2; this was originally believed to be an anti-inflammatory decoy receptor, but is now thought to play a novel and distinct role in sepsis (173). Another C5 cleavage product, C5b, initiates the assembly of the C5b-9 membrane attack complex, which induces lysis of complement-targeted bacteria (103) (Fig. 1).

TLRs comprise a family of pattern recognition receptors named after their similarity to the *Drosophila* Toll protein (99, 106). Their discovery, in the late 1990s, has sparked a resurgent interest in innate immunity. Indeed, the study of TLRs has helped us to appreciate the economical specificity of the innate immune system and that adaptive immunity did not

evolve to replace innate immunity, but rather evolved around it. TLRs are transmembrane glycoproteins comprising an N-terminal leucine-rich repeat domain, a transmembrane region and a C-terminal cytoplasmic signaling domain (76, 84). These receptors are primarily expressed by first-line professional phagocytes (e.g. neutrophils, macrophages and dendritic cells) and are thus strategically located for early recognition of microbial pathogens (1). To date, 10 human TLRs have been identified that generally sense and respond to distinct types of microbial structures (Fig. 2). For instance, TLR3 responds to double-stranded viral RNA, TLR4 responds to enterobacterial lipopolysaccharide, TLR5 responds to bacterial flagellin and TLR9 responds to microbial CpG DNA. TLR2 is unique in that it heterodimerizes with signaling partners (TLR1 or TLR6) for detecting and responding to microbial cell-wall components, such as lipoteichoic acid, lipoproteins, fimbriae, or yeast zymosan (1, 12, 90). TLRs (e.g. TLRs 1, 2, 4, 5 and 6) that are mainly responsible for detecting extracellular microbial structures are expressed on the host cell surface, whereas those (e.g. TLRs 3, 7, 8 and 9) specializing in detecting viral or bacterial nucleic acids are appropriately located intracellularly on endocytic vesicles or organelles (Fig. 2). Following

ligand binding, TLR signaling is triggered upon recruitment of adaptor proteins to the TLR cytoplasmic domains, which help propagate the signals to downstream kinases and transcription factors. This ultimately leads to the induction of immunoregulatory genes that activate or suppress the innate immune and inflammatory response (for a detailed review on TLR signal transduction see refs 118, 119). The presence of both common and selective adaptors, in conjunction with the apparent compartmentalization of the TLRs, allows the induction of individual signaling pathways (for at least some TLRs) in addition to a core TLR response (119). It is thus possible that activation of diverse TLR intracellular pathways, dependent upon different TLR ligand specificities, may allow the host to tailor a response that is appropriate against a given pathogen.

Both complement and TLRs are rapidly activated by most pathogens upon encounter with the host, and common microbial molecules, such as gram-negative bacterial lipopolysaccharide and yeast zymosan, can act both as TLR ligands and complement activators. It is conceivable that the coordination of the early innate response would require a crosstalk between the complement system and the TLR system. In this regard, a systematic analysis of crosstalk in intracellular signaling pathways has revealed that a great number of microbe-induced stimuli converge on a relatively limited number of effector signaling pathways (115). In principle, a molecular crosstalk between complement and TLRs could result in cross-regulation of the two systems, including potential synergistic, or even antagonistic, interactions. These interactions may help enhance host defense or regulate it to prevent excessive inflammatory responses. However, it is also plausible that at least some crosstalk interactions may be instigated by the pathogens themselves for deregulating or modifying the host response in a way that favors their survival. Although this issue has only recently started to be addressed, available evidence indicates bidirectional cooperation between the complement system and the TLR system, because complement regulates TLR activation (67, 183), whereas TLR signaling transmodulates the activity of complement receptors (60, 63).

Periodontitis, associated bacteria and complement / TLR immunity

Periodontal disease is possibly the most common chronic disorder of infectious origin in humans,

resulting in inflammatory destruction of the tooth-supporting tissues (125). The disease is initiated by certain species of subgingival gram-negative anaerobic bacteria that co-exist within dynamic communities of highly organized architecture (31, 158), originally termed 'dental plaque', which predates the more modern term 'biofilm' (46, 108). In fact, seldom would discussions on pathogenic biofilms omit the classical example of dental plaque (25, 88). In periodontal health, the ordered structure of the dental plaque biofilm consists predominately of gram-positive, facultative anaerobic bacteria, although the onset of the disease is associated with a shift to gram-negative anaerobic bacteria, which begin to colonize the subgingival pocket with greater frequencies (157). Using a color-coded system, Socransky and colleagues (158) characterized these microbial communities as red, orange, green, purple and yellow complexes, on the basis of cluster analysis, community ordination and associated disease severity. The presence of high numbers of red complex members such as *Porphyromonas gingivalis*, *Treponema denticola* and *Tannerella forsythia* correlates strongly with periodontal tissue destruction (70, 158). *Prevotella intermedia* and *Fusobacterium nucleatum*, both members of the orange complex, are also associated with various forms of periodontal disease (28, 158, 178).

While the bacteria constitute an essential etiologic factor, it is the host inflammatory reaction to bacterial challenge that primarily mediates periodontal tissue damage (43). This is not to say, however, that the challenge in periodontitis involves simply the issue of controlling the inflammatory response. In a related context, purely anti-inflammatory therapies in sepsis clinical trials have generally failed, even if the initial hyperinflammatory stage was controlled; indeed, many patients would succumb to the infection itself at later stages of the disease (137). Therefore, periodontal and other infection-driven inflammatory diseases should be dealt with in ways that address both infection and inflammation. This, in turn, requires an adequate understanding of both protective and destructive aspects of the host response, and how pathogens may evade the former and contribute to the latter.

There is strong evidence that complement and TLRs form an important link between infection and various local or systemic autoimmune or inflammatory conditions, such as septic shock, ulcerative colitis, rheumatoid arthritis, systemic lupus erythematosus, atherosclerosis, ischemia/reperfusion injury and asthma (3, 16, 24, 135, 172). There is also

evidence to support the involvement of the complement and TLR systems in periodontitis. In this regard the complement is abundant in the gingival crevicular fluid, where it is present at up to 70% of its concentration found in serum (130). Activated complement fragments are abundant in the gingival crevicular fluid of periodontitis patients, whereas they are absent or present in lower concentrations in the crevicular fluid from healthy individuals (7, 17, 147, 148). The functionality of the complement components of the gingival crevicular fluid has been confirmed (26), whereas activated complement fragments have also been detected in the gingival connective tissue (26). Importantly, induction of experimental gingivitis in human volunteers causes progressive elevation of complement cleavage products and correlates with increased accumulation of microbial plaque, clinical inflammation and bleeding on probing (123). These clinical findings suggest a role for the involvement of complement in periodontal pathogenesis. Moreover, *in vitro* mechanistic studies have demonstrated complex interactions between periodontal bacteria and the complement system (61, 104, 127, 130, 156).

In addition to elevated complement activity, the inflamed periodontium is infiltrated by TLR-expressing inflammatory cells, whereas healthy gingivae display significantly lower levels of TLR expression (111, 112, 134). Besides professional inflammatory cells, gingival epithelial cells and fibroblasts also express TLRs and the level of expression correlates with disease activity (94, 134, 162, 170). In terms of function, TLRs (particularly TLR2, and to a much lesser extent TLR4) have been shown to regulate important immune and inflammatory responses to periodontal bacteria *in vivo* and *in vitro* (6, 15, 23, 30, 40, 57, 62, 121, 182).

However, the precise roles, whether protective or destructive, played by the complement and the TLRs in periodontal infection and inflammation are poorly understood. This is partly because these issues have not yet been systematically investigated. Nevertheless, a substantial body of available literature exists, which, if properly synthesized and interpreted, could provide important new insights for future studies. Below, we aim to contribute to this direction by discussing pertinent studies from the oral microbiological and immunological literature. We will focus on major pathogens for which adequate literature exists on their interactions with both the complement and the TLR systems, namely, *P. gingivalis*, *T. denticola*, *T. forsythia* and *P. intermedia*.

***P. gingivalis*: master of subversion**

In principle, a host inflammatory response can become destructive when it is deregulated and its magnitude spirals out of proportion to the microbial threat, or when it is undermined by pathogens, leading to persisting, but ineffective, inflammation in terms of infection control (43, 52, 93, 139). In the context of periodontitis, *P. gingivalis* could be reasonably characterized as a master of subversion, on the basis of the sophisticated sabotage tactics presented below. This gram-negative anaerobic organism expresses an elaborate system of adhesins and proteolytic enzymes (e.g. long and short fimbriae, hemagglutinins, and Arg- and Lys-specific cysteine proteinases known as gingipains), which coordinately enable the pathogen to colonize host tissues and secure critical nutrients (97). Important as these virulence features may be, *P. gingivalis* would probably be unable to establish a chronic infection, unless it could have also evolved ways to evade, undermine, or trick the host immune system. This is lucidly exemplified by its capacity to not only subvert both complement and TLR immunity but, moreover, to exploit crosstalk signaling pathways between complement and TLRs.

Neutralization of complement action

P. gingivalis causes significant inhibition of complement activation, regardless of the initiation pathway involved (classical, lectin, or alternative; Fig. 1), through gingipain-dependent degradation of key complement components, such as C3 (reviewed in refs 129, 155). As a consequence, the deposition of opsonins or of the membrane attack complex on the pathogen surface is suppressed, unless its gingipain activity is ablated by chemical or genetic means (145, 156). All three gingipain enzymes participate in complement inactivation, although the Arg-specific enzymes (HRgpA and RgpB) are more potent in this regard than the Lys-specific gingipain (Kgp) (127). As a further safety measure, the pathogen appears to hijack physiological mechanisms of inhibiting the complement cascade. In this regard, *P. gingivalis* uses its HRgpA to capture the circulating C4b-binding protein on the bacterial cell surface, thereby acquiring the ability to negatively regulate the classical pathway C3 convertase (131).

The findings summarized above are consistent with the observations that *P. gingivalis* is exquisitely resistant to the lytic action of complement (127, 156). Curiously, however, Arg- and Lys-gingipain mutants are as resistant as the wild-type organism upon their exposure to human serum, even though active complement fragments are readily deposited on their bacterial surface (156). These intriguing observations suggest an inherent protective mechanism that is independent of complement inactivation. Indeed, a surface anionic polysaccharide was implicated in this inherent resistance because *P. gingivalis* mutants lacking this structure become readily susceptible to complement-mediated lysis (156). Although this anionic polysaccharide may directly confer resistance, the possibility of an indirect effect may not formally be ruled out. In this context, certain pathogens (e.g. *Helicobacter pylori* and *Escherichia coli*) acquire resistance against complement-induced lysis by expressing molecules that can bind CD59, a host regulatory protein that inhibits the terminal step of membrane attack complex formation (96).

It therefore appears that *P. gingivalis* may be using a number of different reinforcing mechanisms to ensure its survival in the presence of complement. In this regard, because the inhibitory mechanisms of *P. gingivalis* against complement activation are leaky (127), it makes sense that it has also developed inherent resistance against complement-dependent lysis. However, if the surface anionic polysaccharide is sufficient to provide inherent protection, a plausible question is why the pathogen has evolved additional ways to suppress a system that cannot kill it. An interesting interpretation of this is that *P. gingivalis* may have evolved complement inactivation capacity not for its own protection, but for the benefit of other organisms occupying the same subgingival niche. This action may not be as altruistic as it seems; it may actually offer a survival advantage for *P. gingivalis* because it depends on other periodontal bacteria for enhanced colonization and full expression of virulence (81, 88, 126). As *P. gingivalis* is resistant to the lytic action of complement (127, 156), the ability of the complement system to directly offer host protection against this organism is seriously questioned. Nevertheless, it cannot be ruled out that complement activation may indirectly fight this pathogen through the recruitment and activation of phagocytic cells. However, as discussed in a later section (Exploitation of crosstalk interactions between TLRs and complement), *P. gingivalis* may have evolved strategies to diminish or evade its destruction by phagocytes in the presence of complement.

Evasion and subversion of TLRs

Currently available evidence suggests that *P. gingivalis* may have also evolved ways to evade or subvert the TLR system, which senses this organism primarily through TLR2, as shown *in vitro* and *in vivo* (15, 57). However, TLR4 appears to play little or no role in cell activation in response to this oral bacterium (15, 57). These observations appear curious given that *P. gingivalis* is a gram-negative organism which expresses a lipopolysaccharide. However, the organism elegantly utilizes specific lipid A 1- and 4'-phosphatases and a deacylase, which, in concert, generate a tetra-acylated and dephosphorylated lipid A structure (21). This modification renders the lipopolysaccharide molecule biologically inert, thereby allowing *P. gingivalis* to evade TLR4 activation (21). At the same time, this modification confers protection against polymyxin B and perhaps other cationic antimicrobial peptides (21). Intriguingly, the presence of high concentrations of hemin (an environmental nutrient found in diseased sites) suppresses lipid A 1-phosphatase activity and leads to the production of a monophosphorylated lipid A, which actively antagonizes TLR4 activation (21, 22). Thus, even though *P. gingivalis* may express other molecules with intrinsic TLR4 agonistic activity, TLR4 activation is probably suppressed in the context of the whole organism (Fig. 3), as seen both *in vitro* and *in vivo* (15, 57). In this regard, *P. gingivalis* behaves like certain other, nonoral pathogens that have also opted to modify their surface structures in order to escape recognition by TLR4 (8, 109, 152).

The above considerations may explain why TLR2, rather than TLR4, is the predominant TLR involved in recognition of *P. gingivalis*. Induction of TLR2 signaling by *P. gingivalis* requires a signaling partner (TLR1 or TLR6), takes place in membrane lipid rafts (where the receptors are recruited ad hoc) and is facilitated by a nonsignaling coreceptor (CD14) which constitutively resides in lipid rafts (57). Although the host TLR2 response may be potentially protective, *P. gingivalis* has developed ways to undermine the intended host response. Indeed, the pathogen was shown to manipulate the TLR2 response by instigating a molecular crosstalk between TLR2 and the CXC-chemokine receptor 4 in macrophage lipid rafts (62). Specifically, the binding of *P. gingivalis* fimbriae to CXC-chemokine receptor 4 induces cyclic AMP-dependent protein kinase A signaling, which in turn suppresses the TLR2-dependent activation of nuclear factor- κ B and the induction of nitric oxide (Fig. 3) (62). The inhibition of production of this key




Fig. 3. Evasion or subversion of Toll-like receptor (TLR) activation by *Porphyromonas gingivalis*. *P. gingivalis* uses an elaborate system of lipid A phosphatase and deacylase activities that modify the lipid A structure of its lipopolysaccharide (21, 22). These modifications result in lipopolysaccharide molecules that can either evade or actively antagonize TLR4 activation (depicted as a homodimer; TLR4/4) (21, 22). Although the activation of the TLR2/TLR1 heterodimer (TLR2/1) is not antagonized at

the TLR receptor level, *P. gingivalis* instigates a molecular crosstalk between the CXC-chemokine receptor 4 and TLR2/1. Unlike CD14, which facilitates TLR2/1 activation by the pathogen (57), CXCR4 suppresses TLR2 signaling (62). Mechanistically, *P. gingivalis* uses its fimbriae to bind CXCR4 and induce cyclic AMP-dependent protein kinase A (PKA) signaling, which in turn inhibits the activation of nuclear factor-kappaB (NF-κB) activation (62).

antimicrobial molecule promotes the ability of *P. gingivalis* to survive *in vitro* and *in vivo* (62).

The impact of TLR2 signaling on the ability of *P. gingivalis* to cause experimental periodontitis was examined by two independent studies, which found that TLR2-deficient mice (but not TLR4-deficient mice or wild-type controls) are protected against periodontal bone loss (15, 47). These findings are consistent with the notion that TLR2 signaling is manipulated by *P. gingivalis* in a way that promotes its virulence. However, an alternative or additional interpretation is that the observed enhanced bone loss in normal mice could be attributed to *P. gingivalis* induction of TLR2-mediated inflammatory osteoclastogenesis (169).

Exploitation of crosstalk interactions between TLRs and complement

TLR2 activation by *P. gingivalis* induces two distinct signaling cascades (60). One of the cascades leads to

the induction of proinflammatory and antimicrobial responses, and represents the pathway that is manipulated by *P. gingivalis* through exploitation of CXC-chemokine receptor 4, as outlined above. The other cascade represents a pro-adhesive pathway and involves a crosstalk between TLR2 and the complement system (60). Specifically, *P. gingivalis* induces TLR2 inside-out signaling, which transactivates the adhesive capacity of complement receptor-3 (64) (Fig. 4). This crosstalk is made possible by the property of complement receptor-3 to cluster with TLRs in lipid rafts of *P. gingivalis*-stimulated cells (57). Once transactivated, however, complement receptor-3 becomes a target of subversive activity by *P. gingivalis*.

Indeed, *P. gingivalis* uses its fimbriae to bind complement receptor-3, which in turn mediates the uptake of this oral pathogen by macrophages (59). Intriguingly, this phagocytic mechanism does not promote the killing of *P. gingivalis* (171), possibly because complement receptor-3 is not linked to vigorous microbicidal mechanisms (139). In contrast, when *P. gingivalis* is phagocytosed by alternative




Fig. 4. Crosstalk pathways between Toll-like receptors (TLRs) and complement in *Porphyromonas gingivalis*-activated macrophages. TLR recognition of *P. gingivalis* is predominantly mediated by the TLR2/TLR1 heterodimer (TLR2/1), aided by the CD14 co-receptor (57). This interaction induces phosphatidylinositol 3-kinase (PI3K)-dependent inside-out signaling, which transactivates the high-affinity state of complement receptor-3 (CR3) (54, 64) (CR3). Interestingly, *P. gingivalis* interacts with activated CR3 and induces extracellular signal-related kinase 1/2 (ERK1/2) signaling, which in turn downregulates the expression of messenger RNA for cytokines of the inter-

leukin-12 family (56). Moreover, *P. gingivalis* uses its gingipains to attack C5 and release biologically active C5a (127, 176). Through its receptor (C5aR), C5a can activate PI3K and ERK1/2, which in turn suppress critical transcription factors (the interferon regulatory factors 1 and 8; IRF-1 and -8), required for expression of cytokines of the interleukin-12 family (67). Intriguingly, inhibition of bioactive interleukin-12 through these mechanisms results in impaired immune clearance of *P. gingivalis* *in vivo* (56), suggesting that the pathogen exploits TLR/complement crosstalk signaling to promote its virulence.

receptors (i.e. when complement receptor-3 is blocked or genetically ablated), the intracellular killing of this pathogen is dramatically enhanced (171).

The interaction of *P. gingivalis* with complement receptor-3 also activates the extracellular signal-regulated kinase 1/2, which in turn selectively inhibits expression of messenger RNA for the p35 and p40 subunits of interleukin-12 (56) (Fig. 4). Interleukin-12 is a key cytokine involved in pathogen clearance through regulating the production of interferon- γ , which is a potent activator of the macrophage microbicidal capacity (168). Consistent with the above, wild-type mice elicit lower levels of interleukin-12 and interferon-gamma, and display impaired clearance of *P. gingivalis* systemic infection, compared with mice that lack complement receptor-3 (56). Similar results are seen after CR3 blockade with a specific antagonist, which, furthermore, suppresses *P. gingivalis* induction of perio-

dontal bone loss in mice (56). In brief, there is compelling evidence that complement receptor-3 constitutes an Achilles' heel which confers host susceptibility to *P. gingivalis* infection. In this regard, it seems likely that *P. gingivalis* may have actually co-opted a natural anti-inflammatory mechanism to evade innate immunity. Specifically, complement receptor-3 is heavily committed with phagocytosis of iC3b-coated apoptotic cells, which are not normally recognized as danger (83, 107). This precludes induction of a vigorous host response and, in fact, the production of interleukin-12 is inhibited following the phagocytosis of apoptotic cells by macrophages (83).

Although *P. gingivalis* inhibits the complement cascade, curiously enough, the pathogen proactively generates one of the active complement fragments. Specifically, all three gingipains (HRgpA, RgpB and Kgp) act in a C5 convertase-like manner and generate biologically active C5a through the limited degrada-

tion of C5, whereas the C5b remnant is functionally inert (127, 176). When C5 is oxidized by hydroxyl radicals (as may occur in the oxidative environment of the inflammatory response) the gingipains generate increased C5a biological activity (35). Furthermore, *P. gingivalis* may indirectly generate functional C5a by exploiting the physiological crosstalk between the coagulation and the complement systems, which activates the so-called extrinsic pathway (71) (Fig. 1). Indeed, HRgpA and RgpB activate prothrombin to form thrombin (73), which, in turn, generates biologically active C5a by acting as a C5 convertase (71) (Fig. 1). Although C5a can potentially play a key role in host defense against infection (50), it seems highly unlikely that *P. gingivalis* uses its enzymes to generate C5a to contribute to its elimination.

An intriguing question, therefore, is whether there is any selective pressure or advantage for *P. gingivalis* to specifically generate C5a, given that this chronically persisting pathogen overall inhibits the complement cascade. A possible scenario is that the local generation of excessive levels of C5a could incapacitate the antimicrobial function of gingival crevicular neutrophils, rendering them less threatening to *P. gingivalis*. This is because neutrophils become immunologically paralyzed in the presence of high concentrations (10–100 nm) of C5a and thereby fail to carry out functions such as chemotaxis, phagocytosis and production of antimicrobial and inflammatory mediators (72, 172). Such immunological dysfunction has been seen both *in vitro* and *in vivo* and involves both human and rodent neutrophils (72, 136, 172). In fact, C5a-mediated inhibition of neutrophil killing of *P. gingivalis* does occur, at least *in vitro* (J. Krauss and G. Hajishengallis, unpublished data). However, the underlying mechanisms, whether involving immune paralysis or alteration of specific signaling pathways, are currently under investigation. In addition to its potential exploitation by *P. gingivalis*, C5a may amplify periodontal tissue damage through its ability to recruit and activate inflammatory cells. For example, enhanced production of reactive oxygen species by C5a-stimulated neutrophils (50) may contribute to oxidative periodontal tissue destruction (19). On the other hand, this host response would not affect *P. gingivalis*, because it is resistant to killing by reactive oxygen species (58, 113).

Even if the gingipain activity of *P. gingivalis* is capable of increasing the microenvironmental C5a concentrations to paralyzing levels for the neutrophils, this would not impinge on the function of macrophages, which can also be recruited to the

gingival crevice or additionally interact with the pathogen in the periodontal connective tissue (33, 166). Indeed, macrophages are quite resistant to the deleterious effects of high C5a concentrations, because they express relatively modest levels of the C5a receptor relative to the neutrophils (172). For instance, whereas the ability of neutrophils to induce tumor necrosis factor- α (and other innate responses) is inhibited in the presence of C5a at concentrations of ≥ 10 nm, the macrophages display potentiated tumor necrosis factor- α responses under the same C5a concentrations (72, 136).

Therefore, even at high levels, C5a does not exert a general immunosuppressive influence on macrophages. Strikingly, however, C5a can specifically downregulate cytokines of the interleukin-12 family. Indeed, C5a-induced signaling in macrophages interferes with TLR-induced expression of mRNA for the interleukin-12 p35, interleukin-12/interleukin-23 p40 and interleukin-23 p19 subunits (67, 143). These regulatory effects are possibly mediated through C5a-induced phosphatidylinositol 3-kinase and extracellular signal-regulated kinase 1/2 signaling, which in concert suppress critical transcription factors, namely the interferon regulatory factors 1 and 8 (67) (Fig. 4). At the protein level, the production of interleukin-12 is inhibited both *in vitro* and *in vivo*, leading to suppression of T-helper type 1 cell-mediated immunity (67, 183). Moreover, the ability of C5a to inhibit the expression of messenger RNA of both interleukin-23 subunits strongly suggests that C5a can interfere with the capacity of this cytokine to support the development of the T-helper type 17 cell subset (11). The physiological significance of these C5a regulatory effects is likely to attenuate potential tissue damage mediated by T-helper type 1 and T-helper type 17 cells, as seen in various pathological inflammatory conditions (43, 101, 144). However, undesirable outcomes may arise when C5a is not produced physiologically but through the uncontrolled action of microbial enzymes, such as the *P. gingivalis* gingipains. As interleukin-12 is important for immune control of *P. gingivalis* (56), it is possible that this pathogen may exploit the C5a-induced crosstalk with TLR2 for inhibiting IL-12-dependent immune clearance. Such an evasion mechanism may be complementary, rather than redundant, because the interaction of *P. gingivalis* with complement receptor-3 causes partial inhibition (about 60%) of interleukin-12 production (56). The notion that *P. gingivalis* hijacks C5a for its own benefit is additionally supported by observations that the intracellular survival of this pathogen in

macrophages is promoted in the presence of C5a (M. Wang and G. Hajishengallis, unpublished observations).

Interestingly, unlike C5a, C3a is extensively degraded by *P. gingivalis* gingipains and does not retain biological activity (176). Whether this is beneficial for the pathogen is uncertain, but it should be noted that C3a exerts direct antimicrobial effects and readily kills both gram-negative and gram-positive bacteria such as *E. coli*, *Pseudomonas aeruginosa* and *Enterococcus faecalis* (116). If C3a can also kill *P. gingivalis*, then its gingipain-mediated inactivation would serve to protect *P. gingivalis*.

In summary, it appears that *P. gingivalis* does not have a purely defensive agenda in dealing with the complement system. In other words, the pathogen may not restrict its action to simply inhibiting the complement cascade, but rather may proactively employ specific complement components (such as the complement receptor-3 and the C5a) for bidirectional crosstalk interactions with TLR2 that favor the pathogen (Fig. 4). These and other subversive mechanisms (for a review see refs 52, 129, 181) justify the characterization of *P. gingivalis* as a keynote periodontal pathogen, in that it can contribute virulence attributes that are essential for the survival of the entire biofilm community (28, 52).

***T. denticola*: spiraling out of control**

In health, spirochetes (spiral-shaped bacteria with periplasmic flagella) inhabit the periodontal tissues at relatively low frequencies. In sharp contrast, a progressive decline in periodontal health correlates positively with a surge in the spirochete population, which constitutes a substantial percentage of the plaque microbiota in clinically diseased sites (38, 149). Although their role in the initiation of periodontal diseases is uncertain, treponemes are literally at the forefront of established periodontal lesions in dominating numbers (38). In particular, an elevation in the numbers of *T. denticola*, a relatively well-characterized oral spirochete, correlates with progressive disease activity and signifies a defining risk factor for advanced pocketing and periodontal attachment loss (149, 158). Being a member of the red complex, *T. denticola* can engage in specific co-aggregating interactions with the other members (*P. gingivalis* and *T. forsythia*) and with *F. nucleatum*, mediated mostly by its 53-kDa major outer sheath protein (138, 149).

The ability of *T. denticola* to colonize and invade the periodontium can be attributed, at least in part, to its remarkable motility, multiple adhesive properties and high proteolytic activity (18). In this regard, dentilisin, a cell-surface adhesin with chymotrypsin-like properties, appears to facilitate the invasion of connective tissues via the degradation of epithelial cell tight junctions (20). Moreover, dentilisin promotes bacterial invasion of the gingivae by activating matrix metalloproteinases to degrade the extracellular matrix, a mechanism, which at the same time, contributes to periodontal tissue destruction (34, 38). However, the capacity of *T. denticola* to chronically persist in periodontal tissues suggests that the pathogen may have evolved additional ways to inactivate or escape immune-defense mechanisms. In this context, we review below its interactions with the complement and the TLR systems.

Complementary employment of the enemy's arsenal

Because of its potentially destructive effect on host tissues, complement activation is tightly controlled by membrane-bound proteins as well as by soluble regulatory host proteins. One such soluble control protein is factor H, a 155-kDa glycoprotein that controls the alternative pathway by inhibiting the formation of, and accelerating the decay of, the alternative pathway C3 convertase (2). Moreover, factor H contributes to the cleavage and inactivation of C3b to iC3b, and thus can attenuate the activation of the classical pathway C5 convertase and the downstream proinflammatory events (2).

Although the most direct way whereby pathogens can inhibit complement activation involves the use of virulence proteases that degrade and inactivate complement components, another mechanism of evasion depends upon the ability of the pathogen to hijack and employ circulating complement regulators (96). In this regard, *T. denticola* expresses an 11.4-kDa cell-surface lipoprotein that can bind factor H (104). This factor H-binding protein has little or no homology to other bacterial virulence proteins with similar function, suggesting convergent evolution for possible evasion of the complement system. Strikingly, however, once full-length factor H becomes associated with *T. denticola*, the organism uses its serine protease dentilisin to generate a 50-kDa factor H fragment that remains attached to the bacterial surface (104). This seems paradoxical because

dentilisin appears to negate the action of the factor H-binding protein. However, the dentilisin-mediated proteolysis of factor H is quite slow as this complement protein remains essentially intact after 40 min and complete digestion requires 1h (104). Because complement activation is a remarkably rapid process, there is plenty of useful time during which *T. denticola* can escape complement killing by means of surface-bound, biologically active factor H. An interesting question that remains unanswered is whether the attached fragment retains useful complement-regulatory activity or whether it might be employed to serve a different virulence role. What is quite certain, however, is that proactive acquisition of factor H and its proteolytic processing would not be selected if it was detrimental for the pathogen.

The *T. denticola* dentilisin can also use another complement component as substrate, to generate a biologically active fragment. Specifically, this bacterial enzyme generates iC3b upon hydrolysis of the α chain of C3 (179). This is consistent with an early study showing that *T. denticola* is readily opsonized with iC3b (146). As iC3b-opsonized bacteria are taken up by phagocytes via complement receptor-3, it would seem counterproductive for a pathogen to help the host generate even more iC3b. However, as alluded to above, iC3b-mediated phagocytosis is often associated with weak killing mechanisms, or even with immunosuppressive signaling (10, 83, 107, 177), which may be exploited by certain pathogens. Indeed, in addition to *P. gingivalis* discussed above, other microbial pathogens, such as *Mycobacterium tuberculosis*, *Bordetella pertussis* and human immunodeficiency virus-1 (HIV-1), promote their survival by exploiting complement receptor-3-mediated entry, either by direct interaction with the receptor or upon opsonization with iC3b (39, 68, 96).

In summary, although *T. denticola* definitely cannot prevent the initiation of the complement cascade (146), it may be protected from its killing potential by hijacking and employing specific downregulatory factors or specific opsonins for safe intracellular entry. However, additional functional studies are warranted to confirm these plausible mechanisms. Intriguingly, even if *T. denticola* was unable to resist complement-dependent killing successfully, its co-existence with *P. gingivalis* in shared niches (158) may provide it with effective protection mechanisms. *In vivo* observations in mouse models, that mixed infections with both pathogens are more virulent than mono-infections (80, 81, 85), are consistent with this notion, although the precise synergistic mechanisms operating *in vivo* have not been elucidated.

Interactions with TLRs: paying a Toll to escape?

Studies in gingival epithelial cells, macrophages and transfected cell lines have shown that *T. denticola* activates TLR2, rather than TLR4 (5, 13, 117, 141). When purified molecules of *T. denticola* were examined for cell activation, the major outer sheath protein was confirmed to be a TLR2 agonist, although its lipooligosaccharide induced the production of cytokines and nitric oxide in a TLR4-dependent, but TLR2-independent, manner (117). It is uncertain at present whether the lipooligosaccharide is not properly exposed on the bacterial cell surface or if the pathogen possesses TLR4-antagonistic activity, similarly to *P. gingivalis* (22), as a means to evade TLR4 activation. Although macrophages readily elicit cytokine production in response to *T. denticola* (117, 141), gingival epithelial cells stimulated with the same pathogen fail to induce interleukin-8 (5, 13). Whereas degradation of this cytokine may in part explain the findings (5), it appears that the inhibitory mechanism in the epithelial cells is mainly exerted at the transcriptional level (13). As gingival epithelial cell production of interleukin-8 is important for chemoattraction and activation of neutrophils, *T. denticola* appears to be capable of proactively suppressing or delaying the neutrophil influx. Such a mechanism, if indeed operating *in vivo*, would be most important, especially in the initial stages of the infection process (i.e. before a relatively recalcitrant pathogenic biofilm has been established). The capacity to suppress interleukin-8 production, as well as to inhibit induction of the human beta-defensin 2 (13), is consistent with observations that the pathogen is frequently seen in close association with the gingival epithelium (38). Moreover, the *T. denticola* mechanism to suppress TLR2-induced production of interleukin-8 is reminiscent of the so-called 'local chemokine paralysis' mechanism used by *P. gingivalis* (29).

Interestingly, *T. denticola* may use an additional mechanism to prevent the influx of neutrophils into its niche. Specifically, its major outer sheath protein was shown to impair Rac1-dependent neutrophil chemotaxis (102). Rac1 is a small GTPase involved in various cellular functions and has a role in the activation of the nicotinamide adenine dinucleotide phosphate (NADPH) oxidase during the generation of the neutrophil oxidative burst (37). Interestingly, *T. denticola* was shown to suppress the induction of this oxygen-dependent killing mechanism in neu-

trophils (150), which is a TLR-dependent function (142). Whether the underlying mechanism involves the ability of *T. denticola* to inhibit Rac1 activity is uncertain, however. In addition to its roles in chemoattractant-induced directional motility (175) and the oxidative burst (37), Rac1 is also involved in the TLR2-induced activation of nuclear factor- κ B (4). Thus, it would be interesting to determine whether the capacity of *T. denticola* to interfere with this central signaling regulator causes extensive, or even global, inhibition of the neutrophil antimicrobial function.

***T. forsythia* and *P. intermedia* vs. innate immunity: uncharted territories under exploration**

T. forsythia was described, in 1986, as a new species associated with periodontitis and was originally named *Bacteroides forsythus* (165). Now, 23 years later, numerous clinical studies in humans, and animal model studies, have firmly confirmed the prominent role of this bacterium in the initiation and progression of periodontal disease (164). As a member of the red complex (together with *P. gingivalis* and *T. denticola*), *T. forsythia* is considered a major periodontal pathogen. *P. intermedia* (formerly *Bacteroides melaninogenicus* ss. *intermedius* then *Bacteroides intermedius*), an obligatory anaerobic, black-pigmented, gram-negative rod, is frequently associated with adult periodontitis, acute necrotizing ulcerative gingivitis (Trench Mouth) and pregnancy gingivitis (27, 51). This organism is also involved in extra-oral infections, including aspiration pneumonia and intra-abdominal infection (9, 14). *P. intermedia* is a member of the orange complex (158) but can also co-aggregate with *P. gingivalis* and *T. forsythia* (79, 160). Relatively little is known about how *T. forsythia* and *P. intermedia* interact with innate immunity, and only recently have studies been conducted to elucidate their interplay with complement and Toll-like receptors. The two bacteria will be examined together in the following sections.

***T. forsythia* and *P. intermedia* interactions with complement**

Given the importance of *T. forsythia* in periodontitis, it is perplexing that literally nothing is known about how this pathogen interacts with complement.

Extensive search of PubMed, Medline, Web of Science and Scopus databases using the combination of key words *Bacteroides forsythus* / complement and *Tannerella* / complement revealed not even a single publication. As *T. forsythia* is exposed to complement in gingival crevicular fluid it can only be speculated that, like other major periodontopathic bacteria, the bacterium is either resistant to complement, or it is protected from the microbicidal activity of complement by other inhabitants of dental plaque (see below). Indeed, preliminary data strongly suggest that akin to other periodontopathogens, *T. forsythia* proteases may also play an essential role in neutralizing complement (M. Bojko, J. Potempa, K. Abdulkarim and A. Blom, unpublished results).

Several lines of evidence from early studies suggest that complement is a potentially important host defense mechanism against *P. intermedia*. First, heat inactivation of complement in normal human serum significantly reduces opsonic activity for *P. intermedia* (167). Second, in the absence of the classical pathway, *P. intermedia* is opsonized by the alternative pathway, probably in response to its lipopolysaccharide (69, 120). Nevertheless, the classical pathway is also important because kinetic studies have revealed that opsonization proceeds at significantly faster rates when the classical pathway is intact (167). Third, different strains of *P. intermedia* show variable sensitivity to the complement-dependent bactericidal activity of serum (161). Finally, the alternative pathway was shown to contribute to the killing of serum-sensitive strains, while the classical pathway was primarily responsible for killing strains with intermediate sensitivity (161).

Interestingly, a number of *Prevotella* subspecies and strains, including *P. intermedia*, which have been associated with gingivitis and periodontitis, display higher proteolytic activity than that recorded for commensal strains isolated from healthy mouths (180). This suggests that, akin to the gingipains of *P. gingivalis*, proteases produced by *P. intermedia* may also contribute to resistance against the antibacterial activity of complement. Indeed, recent and detailed investigations clearly implicate interpain A (InpA), a streptopain-like (SpeB-like) cysteine protease secreted by some *P. intermedia* strains, in conferring 100% resistance to killing by 20% normal human serum (130). By contrast, an isogenic interpain A-deficient strain showed a survival rate of only 78% (130). This protective effect is exerted by efficient degradation of complement factor C3, which is the central component of the entire complement system (Fig. 1). Significantly, proteolytic inactivation

of C3 by interpain A occurs in whole human serum, which explains the observation that opsonization and/or opsonization-dependent phagocytosis of *E. coli* and *Proteus mirabilis* are impaired when the serum is pretreated with certain *Bacteroides* species, including *P. intermedia* (77, 114, 167).

Moreover, it appears that *P. intermedia* has developed additional, protease-independent strategies for evading killing by complement. This can be inferred from its significant survival rate, even when the bacterium is incubated with serum containing the broad-spectrum cysteine protease inhibitor, E-64 (128). This inhibitor should ameliorate the proteolytic activity not only of interpain A, but also of two other streptopain-like proteases expressed by *P. intermedia* (128). Although it cannot be excluded that serine proteinase activity, insensitive to E-64 inhibition (151), may be employed by *P. intermedia* to neutralize complement, it seems more likely that other mechanisms, such as binding of host-derived complement inhibitors, or the molecular composition of the bacterial surface, may confer exceptional resistance against complement-dependent lysis.

Oral bacterial co-aggregation is important for dental plaque formation, and *P. intermedia* has been shown to co-aggregate with *P. gingivalis* (78, 79). In the context of this specific interaction and the general pathogenicity of biofilms, it becomes important to note that the interpain of *P. intermedia* synergizes with the gingipains of *P. gingivalis* in complement inactivation (130). It is plausible that secreted interpain and gingipains aid in the survival of bystander bacterial species in dental plaque, which could otherwise be eliminated by the bactericidal activity of complement. This bacterial synergy may be one of the mechanisms that contribute to a favorable biofilm environment which could foster the survival of the entire microbial community.

In interacting with complement, *P. intermedia* shares one important feature with *P. gingivalis*. At low concentrations, the interpain, similarly to gingipains, is able to activate the C1 complex in serum, causing deposition of C1q on bacterial surfaces (130). Furthermore, the interpain can directly release anaphylatoxin C3a, while gingipains additionally release C5a (127, 130). Significantly, although high concentrations of these proteases inhibit the complement cascade, the anaphylatoxins are still released by direct protease action. Taken together, the following scenario can be envisioned (Fig. 5). *P. gingivalis* and *P. intermedia* activate complement when present at low numbers, resulting in a local inflammatory reaction that does not eliminate them because these

bacteria are resistant to complement-mediated lysis. However, complement activation can suppress complement-sensitive bacteria, which could otherwise compete for space and nutrients. The inflammatory serum exudate, moreover, provides *P. gingivalis* and *P. intermedia* with precious nutrients (such as hemin, a source of essential iron). At later stages of infection, the concentration of proteases is high enough to destroy C3 and inhibit complement activation, thus promoting the survival of the entire biofilm community by helping bystander bacteria evade complement killing. Even under these conditions that suppress the physiological activation of the complement cascade, the proactive release of biologically active C5a can stimulate inflammatory serum exudate for acquisition of hemin and other nutrients. Furthermore, because anaphylatoxins are potent mediators of inflammatory reactions (91), their local generation at sites heavily populated with bacteria can contribute to the pathological processes associated with periodontal tissue destruction. Moreover, C5a could be additionally exploited for immune evasion (see above, *Exploitation of crosstalk interactions between TLRs and complement*), thus further promoting pathogen survival and ensuing inflammation. It is thus becoming apparent that periodontal pathogens have evolved in ways that allow them to not only endure inflammation but exploit it for promoting their adaptive fitness, as long as they can shut down critical host responses [such as induction of nitric oxide production (62)] that could eliminate them.

***P. intermedia* and *T. forsythia* interactions with TLRs**

TLR-mediated recognition of *P. intermedia* and *T. forsythia*, and the subsequent activation of signal transduction pathways culminating in the generation of inflammatory cytokines, can have a direct bearing on periodontal pathogenesis. Unfortunately, our knowledge about TLR interactions with these two pathogens is limited. Similarly to *P. gingivalis* and *T. denticola*, which predominantly activate TLR2, *P. intermedia* and *T. forsythia* sonicates exclusively stimulate TLR2 in human cell lines transfected with various combinations of TLRs and co-receptors (82). Consistent with these results, purified lipoproteins of *T. forsythia* and a nonendotoxic glycoprotein of *P. intermedia* activate TLR2 signaling in human monocytic THP-1 cells and/or human gingival fibroblasts (65, 159). By contrast, highly purified lipid




Fig. 5. Biphasic virulence effects of *Porphyromonas gingivalis* and *Prevotella intermedia* cysteine proteases. (A) Effects of low concentrations of the proteases (i.e. early in the colonization process), which activate the classical pathway. (B) Effects of high concentrations of the proteases (i.e. in a developed biofilm), which suppress the physiological activation of the complement cascade, but selectively generate anaphylatoxins via limited degradation of C3 and C5. *P. gingivalis* and *P. intermedia* are exceptionally resistant to the bactericidal activity of complement, and released proteases, gingipains and interpain, respectively, contribute to the resistance. At low concentrations of the proteases, likely to occur at the early stages of infection or at a long distance from the dental

plaque, the released proteases activate the C1 complex, leading to deposition of C1q on the bacterial surface (127, 130). Complement activation may eliminate complement-sensitive commensal bacteria, which could otherwise compete with pathogens for space and nutrients. At high concentrations, the proteases synergistically inhibit the bactericidal activity of complement by degrading C3 and C5, thus protecting complement-sensitive bacteria in their proximity and promoting biofilm development. At the same time, released anaphylatoxins (C3a and C5a; although only the latter was confirmed to retain its biological activity) fuel inflammation, resulting in tissue damage and nutrient generation, and are exploited for immune evasion (see also Fig. 4).

A, derived from the lipopolysaccharide of *P. intermedia*, activates murine peritoneal macrophages to secrete interleukin-6 through a TLR4-dependent pathway (66). However, the fact that whole cells of *P. intermedia* fail to activate TLR4 (82) suggests the possible presence of TLR4 antagonistic activity associated with the bacterial cell surface, although this possibility has not been addressed.

In addition to lipoproteins, another cell surface-associated and secreted protein, the BspA of *T. forsythia*, also activates TLR2, resulting in the

production of interleukin-1, interleukin-8 and tumor necrosis factor-alpha upon interactions with human gingival epithelial cells or human monocytic THP-1 cells (55, 121). In the former interaction, TLR1 serves as a signaling partner (121), whereas the latter interaction is additionally dependent on the CD14 co-receptor (55). From a pathological point of view, it is important to note that *T. forsythia*, *T. denticola* and *P. gingivalis* can induce strongly synergistic interleukin-6 production (163). As interleukin-6 activates osteoclasts (74), this synergistic

Table 1. Exploitation of complement, Toll-like receptors, or their crosstalk by oral pathogens

Mechanism	Pathogen	Effector molecule	References		
1	a. Inhibition of complement activation through digestion of the central complement component (C3)	<i>P. gingivalis</i>	Gingipains, especially HRgpA and RgpB	127	
		<i>P. intermedia</i>	Interpain (InpA)	130	
	b. Synergy in complement inactivation	<i>P. gingivalis</i> & <i>P. intermedia</i>	Gingipains and interpain	130	
2	Inherent resistance to complement-mediated lysis	<i>P. gingivalis</i>	Surface anionic polysaccharide	156	
3	Hijacking complement regulatory proteins	a. C4b-binding protein	<i>P. gingivalis</i>	HrgpA	131
		b. Factor H	<i>T. denticola</i>	11.4-kDa factor H-binding lipoprotein	104
4	Generation of specific complement fragments	a. iC3b (promotes phagocytosis linked to poor microbicidal activity)	<i>T. denticola</i>	Dentilisin	96, 179
		b. C5a (inhibits TLR-induced interleukin-12)	<i>P. gingivalis</i>	HRgpA, RgpB and Kgp	67, 127, 176, 183
5	Promotion of intracellular survival via complement receptor 3-mediated entry	<i>P. gingivalis</i>	Fimbriae	171	
6	TLR4 evasion by expressing dephosphorylated and tetra-acylated lipid A	<i>P. gingivalis</i>	Lipid A 1- and 4'-phosphatases and deacylase	21	
7	TLR4 antagonism by expressing monophosphorylated tetra-acylated lipid A	<i>P. gingivalis</i>	Lipid A 4'-phosphatase and deacylase (lipid A 1-phosphatase suppressed by hemin)	21, 22	
8	Suppression of TLR2-induced interleukin-12 via complement receptor-3 binding	<i>P. gingivalis</i>	Fimbriae	56	
9	Suppression of TLR2 activation through instigated crosstalk with the CXC-chemokine receptor 4	<i>P. gingivalis</i>	Fimbriae	62	
10	Inhibition of TLR2-induced interleukin-8 in gingival epithelial cells	<i>P. gingivalis</i> <i>T. denticola</i>	Unspecified; requires the use of whole cells	13, 29	
11	Suppression of the neutrophil oxidative burst (TLR-dependent)	<i>T. denticola</i>	Unspecified; inhibitory factor released into the culture medium	142, 149	

P. gingivalis, *Porphyromonas gingivalis*; *T. denticola*, *Treponema denticola*.

action may significantly reduce the number of bacteria required to stimulate alveolar bone resorption. This concept becomes biologically relevant in the light of recent evidence that the three pathogens synergize in inducing periodontal bone resorption *in vivo* (81).

Cumulatively, a large body of evidence indicates that periodontal bacteria preferentially activate immune and nonimmune cells using the TLR2 pathway (57, 82, 117, 141, 182). TLR2 stimulation in the absence of simultaneous signaling through TLR4 favors the development of the T helper 2-type immune response (48, 132, 133). Interestingly, the immune response in chronic periodontitis is strongly T helper 2-type biased and is strongly associated with the progressive lesion of periodontitis (44, 45). Therefore, it is reasonable to hypothesize that preferential stimulation of TLR2 by periodontal pathogens contributes to the T helper 2-type skewing of the immune response, which may perpetuate chronic inflammation and periodontal disease progression, as opposed to the T helper 1-type immune response, which is more effective in bacterial clearance by professional phagocytes (43). Additional reasons for this TLR2 ‘preference’ may be related to observations that a number of pathogens, including *P. gingivalis* (see above), have evolved mechanisms for successfully dealing with the outcome of TLR2-dependent cell activation (62, 122, 124, 154, 174). Although there is no compelling evidence that the TLR4 response is necessarily more protective against infection relative to TLR2, it is interesting to note that TLR4, but not TLR2 (as either a TLR2/1 or TLR2/6 heterodimer), activates autophagy in a way that eliminates phagocytosed pathogens by macrophages (32). A notable known exception among oral bacteria that preferentially activate TLR2 is *Aggregatibacter actinomycetemcomitans*, which readily activates TLR4 (82, 182). However, unlike bacteria such as *P. gingivalis*, *T. denticola*, and *P. intermedia*, which may survive by suppressing selective aspects of the host response, *A. actinomycetemcomitans* appears to promote its persistence by elaborating powerful toxins (e.g. leukotoxin and cytolethal distending toxin), which are capable of killing both innate and adaptive immune effector cells (reviewed in ref. 87).

Conclusions and future directions

As discussed above, complement and TLRs function as major defense systems against microbial infection,

whether acting separately or in concert (103, 105, 183). However, in the course of evolution, successful pathogens have ‘learned’ to breach these systems and exploit their communication hubs (96, 140). Table 1 summarizes putative virulence mechanisms whereby periodontal pathogens may resist or escape complement- and TLR-mediated immunity. These microbial strategies may explain, at least in part, the ability of periodontal bacteria, such as *P. gingivalis*, *T. denticola*, *T. forsythia* and *P. intermedia*, to persist and establish chronic infections in the otherwise hostile environment of their host. As these bacteria reside in mixed-species biofilms, it is reasonable to assume a synergy of shared, communal mechanisms of immune subversion to maximize the survival potential of the whole community, a concept that is supported by evidence initially available. This also suggests that even bacteria lacking protective mechanisms on their own may thrive in such environment, as long as their presence is beneficial to the immune-subversive bacteria. For example, although no immune-evasion mechanisms have been reported for *F. nucleatum*, this bacterium is a useful member of the tooth-associated biofilm by forming a co-aggregating bridge between early and late colonizers (92), and can chronically persist in deep periodontal pockets (158).

The analysis of pertinent literature in this review also reveals gaps, especially at the mechanistic level, in our understanding of periodontal bacterial immune evasion. In addition to molecular mechanistic studies that may help to identify novel virulence factors, *in vivo* studies are also warranted to substantiate some of the *in vitro* findings. Although this review has focused on the host defense potential of the complement and the TLRs, and how this is undermined by periodontal bacteria, it should also be noted that both systems contribute to immunopathology in several inflammatory diseases (100, 135). However, relatively little is known regarding the precise roles of the complement and the TLRs in periodontal pathogenesis. Consequently, it is currently uncertain which specific signaling pathways need to be blocked to attenuate the pathology or, conversely, be enhanced to promote host defense. At a first stage, these gaps in our knowledge necessitate mechanistic and interventional studies using appropriate animal models, because causal mechanistic relationships cannot normally be addressed in human studies for important ethical considerations (49). Successful elucidation of the roles of complement and TLRs in immune and inflammatory responses in the periodontium, and how these are proactively modified by peri-

odontal pathogens, will offer insights into periodontal pathogenesis and facilitate the rational design of therapeutic interventions.

Acknowledgment

The authors are supported by grants from the US National Institutes of Health, DE015254, DE018292, DE017138, DE009761, GM062134, AI030040, AI072106 and AI068730.

References

- Akira S, Takeda K. Toll-like receptor signalling. *Nat Rev Immunol* 2004; **4**: 499–511.
- Alexander JJ, Quigg RJ. The simple design of complement factor H: looks can be deceiving. *Mol Immunol* 2007; **44**: 123–132.
- Andreaskos E, Foxwell B, Feldmann M. Is targeting Toll-like receptors and their signaling pathway a useful therapeutic approach to modulating cytokine-driven inflammation? *Immunol Rev* 2004; **202**: 250–265.
- Arbibe L, Mira JP, Teusch N, Kline L, Guha M, Mackman N, Godowski PJ, Ulevitch RJ, Knaus UG. Toll-like receptor 2-mediated NF- κ B activation requires a Rac1-dependent pathway. *Nat Immunol* 2000; **1**: 533–540.
- Asai Y, Jinno T, Ogawa T. Oral treponemes and their outer membrane extracts activate human gingival epithelial cells through toll-like receptor 2. *Infect Immun* 2003; **71**: 717–725.
- Asai Y, Ohyama Y, Gen K, Ogawa T. Bacterial fimbriae and their peptides activate human gingival epithelial cells through Toll-like receptor 2. *Infect Immun* 2001; **69**: 7387–7395.
- Attstrom R, Laurel AB, Lahsson U, Sjöholm A. Complement factors in gingival crevice material from healthy and inflamed gingiva in humans. *J Periodontol Res* 1975; **10**: 19–27.
- Backhed F, Normark S, Schweda EK, Oscarson S, Richter-Dahlfors A. Structural requirements for TLR4-mediated LPS signalling: a biological role for LPS modifications. *Microbes Infect* 2003; **5**: 1057–1063.
- Bartlett JG. Anaerobic bacterial infections of the lung and pleural space. *Clin Infect Dis* 1993; **16** (Suppl. 4): S248–S255.
- Berton G, Laudanna C, Sorio C, Rossi F. Generation of signals activating neutrophil functions by leukocyte integrins: LFA-1 and gp150/95, but not CR3, are able to stimulate the respiratory burst of human neutrophils. *J Cell Biol* 1992; **116**: 1007–1017.
- Bettelli E, Korn T, Kuchroo VK. Th17: the third member of the effector T cell trilogy. *Curr Opin Immunol* 2007; **19**: 652–657.
- Beutler B, Jiang Z, Georgel P, Crozat K, Croker B, Rutschmann S, Du X, Hoebe K. Genetic analysis of host resistance: Toll-like receptor signaling and immunity at large. *Annu Rev Immunol* 2006; **24**: 353–389.
- Brisette CA, Pham TT, Coats SR, Darveau RP, Lukehart SA. *Treponema denticola* does not induce production of common innate immune mediators from primary gingival epithelial cells. *Oral Microbiol Immunol* 2008; **23**: 474–481.
- Brook I. Microbiology of polymicrobial abscesses and implications for therapy. *J Antimicrob Chemother* 2002; **50**: 805–810.
- Burns E, Bachrach G, Shapira L, Nussbaum G. Cutting Edge: TLR2 is required for the innate response to *Porphyromonas gingivalis*: activation leads to bacterial persistence and TLR2 deficiency attenuates induced alveolar bone resorption. *J Immunol* 2006; **177**: 8296–8300.
- Cario E, Podolsky DK. Toll-like receptor signaling and its relevance to intestinal inflammation. *Ann N Y Acad Sci* 2006; **1072**: 332–338.
- Challacombe SJ, Shirlaw PJ. Immunology of diseases of the oral cavity. In: Mestecky J, Lamm ME, Strober W, Bienenstock J, McGhee JR, Mayer L, editors. *Mucosal Immunology*. Amsterdam: Elsevier Academic Press, 2005: 1517–1546.
- Chan EC, McLaughlin R. Taxonomy and virulence of oral spirochetes. *Oral Microbiol Immunol* 2000; **15**: 1–9.
- Chapple IL, Matthews JB. The role of reactive oxygen and antioxidant species in periodontal tissue destruction. *Periodontol 2000* 2007; **43**: 160–232.
- Chi B, Qi M, Kuramitsu HK. Role of dentilisin in *Treponema denticola* epithelial cell layer penetration. *Res Microbiol* 2003; **154**: 637–643.
- Coats SR, Jones JW, Do CT, Braham PH, Bainbridge BW, To TT, Goodlett DR, Ernst RK, Darveau RP. Human Toll-like receptor 4 responses to *P. gingivalis* are regulated by lipid A 1- and 4'-phosphatase activities. *Cell Microbiol* 2009; Epub ahead of print: doi: 10.1111/j.1462-5822.2009.01349.x.
- Coats SR, Pham TT, Bainbridge BW, Reife RA, Darveau RP. MD-2 mediates the ability of tetra-acylated and penta-acylated lipopolysaccharides to antagonize *Escherichia coli* lipopolysaccharide at the TLR4 signaling complex. *J Immunol* 2005; **175**: 4490–4498.
- Coats SR, Reife RA, Bainbridge BW, Pham TT, Darveau RP. *Porphyromonas gingivalis* lipopolysaccharide antagonizes *Escherichia coli* lipopolysaccharide at toll-like receptor 4 in human endothelial cells. *Infect Immun* 2003; **71**: 6799–6807.
- Cook DN, Pisetsky DS, Schwartz DA. Toll-like receptors in the pathogenesis of human disease. *Nat Immunol* 2004; **5**: 975–979.
- Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of persistent infections. *Science* 1999; **284**: 1318–1322.
- Courts FJ, Boackle RJ, Fudenberg HH, Silverman MS. Detection of functional complement components in gingival crevicular fluid from humans with periodontal diseases. *J Dent Res* 1977; **56**: 327–331.
- Darby I, Curtis M. Microbiology of periodontal disease in children and young adults. *Periodontol 2000* 2001; **26**: 33–53.
- Darveau RP. The oral microbial consortium's interaction with the periodontal innate defense system. *DNA Cell Biol* 2009; **28**: 389–395.

29. Darveau RP, Belton CM, Reife RA, Lamont RJ. Local chemokine paralysis, a novel pathogenic mechanism for *Porphyromonas gingivalis*. *Infect Immun* 1998; **66**: 1660–1665.
30. Darveau RP, Pham TT, Lemley K, Reife RA, Bainbridge BW, Coats SR, Howald WN, Way SS, Hajjar AM. *Porphyromonas gingivalis* lipopolysaccharide contains multiple lipid A species that functionally interact with both toll-like receptors 2 and 4. *Infect Immun* 2004; **72**: 5041–5051.
31. Davey ME, Costerton JW. Molecular genetics analyses of biofilm formation in oral isolates. *Periodontol 2000* 2006; **42**: 13–26.
32. Delgado MA, Elmaoued RA, Davis AS, Kyei G, Deretic V. Toll-like receptors control autophagy. *EMBO J* 2008; **27**: 1110–1121.
33. Delima AJ, Van Dyke TE. Origin and function of the cellular components in gingival crevice fluid. *Periodontol 2000* 2003; **31**: 55–76.
34. Ding Y, Uitto VJ, Haapasalo M, Lounatmaa K, Konttinen YT, Salo T, Grenier D, Sorsa T. Membrane components of *Treponema denticola* trigger proteinase release from human polymorphonuclear leukocytes. *J Dent Res* 1996; **75**: 1986–1993.
35. Discipio RG, Daffern PJ, Kawahara M, Pike R, Travis J, Hugli TE, Potempa J. Cleavage of human complement component C5 by cysteine proteinases from *Porphyromonas (Bacteroides) gingivalis*. Prior oxidation of C5 augments proteinase digestion of C5. *Immunology* 1996; **87**: 660–667.
36. Dixon DR, Bainbridge BW, Darveau RP. Modulation of the innate immune response within the periodontium. *Periodontol 2000* 2004; **35**: 53–74.
37. El-Benna J, Dang PM, Gougerot-Pocidalo MA. Priming of the neutrophil NADPH oxidase activation: role of p47phox phosphorylation and NOX2 mobilization to the plasma membrane. *Semin Immunopathol* 2008; **30**: 279–289.
38. Ellen RP, Galimanas VB. Spirochetes at the forefront of periodontal infections. *Periodontol 2000* 2005; **38**: 13–32.
39. Ernst JD. Macrophage receptors for *Mycobacterium tuberculosis*. *Infect Immun* 1998; **66**: 1277–1281.
40. Eskan MA, Hajishengallis G, Kinane DF. Differential activation of human gingival epithelial cells and monocytes by *Porphyromonas gingivalis* fimbriae. *Infect Immun* 2007; **75**: 892–898.
41. Fearon DT. Seeking wisdom in innate immunity. *Nature* 1997; **388**: 323–324.
42. Finlay BB, McFadden G. Anti-immunology: evasion of the host immune system by bacterial and viral pathogens. *Cell* 2006; **124**: 767–782.
43. Gaffen SL, Hajishengallis G. A new inflammatory cytokine on the block: re-thinking periodontal disease and the Th1/Th2 paradigm in the context of Th17 cells and IL-17. *J Dent Res* 2008; **87**: 817–828.
44. Gemmell E, Yamazaki K, Seymour GJ. Destructive periodontitis lesions are determined by the nature of the lymphocytic response. *Crit Rev Oral Biol Med* 2002; **13**: 17–34.
45. Gemmell E, Yamazaki K, Seymour GJ. The role of T cells in periodontal disease: homeostasis and autoimmunity. *Periodontol 2000* 2007; **43**: 14–40.
46. Gibbons RJ, Socransky SS, Dearaujo WC, Vanhoute J. Studies of the predominant cultivable microbiota of dental plaque. *Arch Oral Biol* 1964; **11**: 365–370.
47. Gibson FC 3rd, Ukai T, Genco CA. Engagement of specific innate immune signaling pathways during *Porphyromonas gingivalis* induced chronic inflammation and atherosclerosis. *Front Biosci* 2008; **13**: 2041–2059.
48. Goriely S, Neurath MF, Goldman M. How microorganisms tip the balance between interleukin-12 family members. *Nat Rev Immunol* 2008; **8**: 81–86.
49. Graves DT, Fine D, Teng Y-TA, Van Dyke TE, Hajishengallis G. The use of rodent models to investigate host-bacteria interactions related to periodontal diseases. *J Clin Periodontol* 2008; **35**: 89–105.
50. Guo RF, Ward PA. Role of C5a in inflammatory responses. *Annu Rev Immunol* 2005; **23**: 821–852.
51. Gursoy M, Haraldsson G, Hyvonen M, Sorsa T, Pajukanta R, Kononen E. Does the frequency of *Prevotella intermedia* increase during pregnancy? *Oral Microbiol Immunol* 2009; **24**: 299–303.
52. Hajishengallis G. *Porphyromonas gingivalis*-host interactions: open war or intelligent guerilla tactics? *Microbes Infect* 2009; **11**: 637–645.
53. Hajishengallis G. Toll gates to periodontal host modulation and vaccine therapy. *Periodontol 2000* 2009; **51**: 181–207.
54. Hajishengallis G, Harokopakis E. *Porphyromonas gingivalis* interactions with complement receptor 3 (CR3): innate immunity or immune evasion? *Front Biosci* 2007; **12**: 4547–4557.
55. Hajishengallis G, Martin M, Sojar HT, Sharma A, Schifferle RE, DeNardin E, Russell MW, Genco RJ. Dependence of bacterial protein adhesins on Toll-like receptors for proinflammatory cytokine induction. *Clin Diagn Lab Immunol* 2002; **9**: 403–411.
56. Hajishengallis G, Shakhathreh M-AK, Wang M, Liang S. Complement receptor 3 blockade promotes IL-12-mediated clearance of *Porphyromonas gingivalis* and negates its virulence *in vivo*. *J Immunol* 2007; **179**: 2359–2367.
57. Hajishengallis G, Tapping RI, Harokopakis E, Nishiyama S-I, Ratti P, Schifferle RE, Lyle EA, Triantafilou M, Triantafilou K, Yoshimura F. Differential interactions of fimbriae and lipopolysaccharide from *Porphyromonas gingivalis* with the Toll-like receptor 2-centred pattern recognition apparatus. *Cell Microbiol* 2006; **8**: 1557–1570.
58. Hajishengallis G, Wang M, Bagby GJ, Nelson S. Importance of TLR2 in early innate immune response to acute pulmonary infection with *Porphyromonas gingivalis* in mice. *J Immunol* 2008; **181**: 4141–4149.
59. Hajishengallis G, Wang M, Harokopakis E, Triantafilou M, Triantafilou K. *Porphyromonas gingivalis* fimbriae proactively modulate β_2 integrin adhesive activity and promote binding to and internalization by macrophages. *Infect Immun* 2006; **74**: 5658–5666.
60. Hajishengallis G, Wang M, Liang S. Induction of distinct TLR2-mediated proinflammatory and proadhesive signaling pathways in response to *Porphyromonas gingivalis* fimbriae. *J Immunol* 2009; **182**: 6690–6696.
61. Hajishengallis G, Wang M, Liang S, Shakhathreh MA, James D, Nishiyama S, Yoshimura F, Demuth DR. Subversion of innate immunity by periodontopathic bacteria via

- exploitation of complement receptor-3. *Adv Exp Med Biol* 2008; **632**: 203–219.
62. Hajishengallis G, Wang M, Liang S, Triantafilou M, Triantafilou K. Pathogen induction of CXCR4/TLR2 cross-talk impairs host defense function. *Proc Natl Acad Sci USA* 2008; **105**: 13532–13537.
 63. Harokopakis E, Albzreh MH, Martin MH, Hajishengallis G. TLR2 transmodulates monocyte adhesion and transmigration via Rac1- and PI3K-mediated inside-out signaling in response to *Porphyromonas gingivalis* fimbriae. *J Immunol* 2006; **176**: 7645–7656.
 64. Harokopakis E, Hajishengallis G. Integrin activation by bacterial fimbriae through a pathway involving CD14, Toll-like receptor 2, and phosphatidylinositol-3-kinase. *Eur J Immunol* 2005; **35**: 1201–1210.
 65. Hasebe A, Yoshimura A, Into T, Kataoka H, Tanaka S, Arakawa S, Ishikura H, Golenbock DT, Sugaya T, Tsuchida N, Kawanami M, Hara Y, Shibata K. Biological activities of *Bacteroides forsythus* lipoproteins and their possible pathological roles in periodontal disease. *Infect Immun* 2004; **72**: 1318–1325.
 66. Hashimoto M, Asai Y, Tamai R, Jinno T, Umatani K, Ogawa T. Chemical structure and immunobiological activity of lipid A from *Prevotella intermedia* ATCC 25611 lipopolysaccharide. *FEBS Lett* 2003; **543**: 98–102.
 67. Hawlisch H, Belkaid Y, Baelder R, Hildeman D, Gerard C, Kohl J. C5a negatively regulates toll-like receptor 4-induced immune responses. *Immunity* 2005; **22**: 415–426.
 68. Hellwig SM, van Oirschot HF, Hazenbos WL, van Spriel AB, Mooi FR, van De Winkel JG. Targeting to Fcγ receptors, but not CR3 (CD11b/CD18), increases clearance of *Bordetella pertussis*. *J Infect Dis* 2001; **183**: 871–879.
 69. Hofstad T. Serological properties of lipopolysaccharide from oral stains of *Bacteroides melaninogenicus*. *J Bacteriol* 1969; **97**: 1078–1082.
 70. Holt SC, Ebersole JL. *Porphyromonas gingivalis*, *Treponema denticola*, and *Tannerella forsythia*: the “red complex”, a prototype polybacterial pathogenic consortium in periodontitis. *Periodontol 2000* 2005; **38**: 72–122.
 71. Huber-Lang M, Sarma JV, Zetoune FS, Rittirsch D, Neff TA, McGuire SR, Lambris JD, Warner RL, Flierl MA, Hoesel LM, Gebhard F, Younger JG, Drouin SM, Wetsel RA, Ward PA. Generation of C5a in the absence of C3: a new complement activation pathway. *Nat Med* 2006; **12**: 682–687.
 72. Huber-Lang MS, Younkin EM, Sarma JV, McGuire SR, Lu KT, Guo RF, Padgaonkar VA, Curnutte JT, Erickson R, Ward PA. Complement-induced impairment of innate immunity during sepsis. *J Immunol* 2002; **169**: 3223–3231.
 73. Imamura T, Banbula A, Pereira PJ, Travis J, Potempa J. Activation of human prothrombin by arginine-specific cysteine proteinases (Gingipains R) from *Porphyromonas gingivalis*. *J Biol Chem* 2001; **276**: 18984–18991.
 74. Ishimi Y, Miyaura C, Jin CH, Akatsu T, Abe E, Nakamura Y, Yamaguchi A, Yoshiki S, Matsuda T, Hirano T, Kishimoto T, Suda T. IL-6 is produced by osteoblasts and induces bone resorption. *J Immunol* 1990; **145**: 3297–3303.
 75. Janeway CA Jr, Medzhitov R. Innate immune recognition. *Annu Rev Immunol* 2002; **20**: 197–216.
 76. Jin MS, Kim SE, Heo JY, Lee ME, Kim HM, Paik SG, Lee H, Lee JO. Crystal structure of the TLR1-TLR2 heterodimer induced by binding of a tri-acylated lipopeptide. *Cell* 2007; **130**: 1071–1082.
 77. Jones GR, Gemmell CG. Impairment by *Bacteroides* species of opsonisation and phagocytosis of enterobacteria. *J Med Microbiol* 1982; **15**: 351–361.
 78. Kamaguchi A, Nakayama K, Ohyama T, Watanabe T, Okamoto M, Baba H. Coaggregation of *Porphyromonas gingivalis* and *Prevotella intermedia*. *Microbiol Immunol* 2001; **45**: 649–656.
 79. Kamaguchi A, Ohyama T, Sakai E, Nakamura R, Watanabe T, Baba H, Nakayama K. Adhesins encoded by the gingipain genes of *Porphyromonas gingivalis* are responsible for co-aggregation with *Prevotella intermedia*. *Microbiology* 2003; **149**: 1257–1264.
 80. Kesavalu L, Holt SC, Ebersole JL. Virulence of a polymicrobial complex, *Treponema denticola* and *Porphyromonas gingivalis*, in a murine model. *Oral Microbiol Immunol* 1998; **13**: 373–377.
 81. Kesavalu L, Sathishkumar S, Bakthavatchalu V, Matthews C, Dawson D, Steffen M, Ebersole JL. Rat model of polymicrobial infection, immunity, and alveolar bone resorption in periodontal disease. *Infect Immun* 2007; **75**: 1704–1712.
 82. Kikkert R, Laine ML, Aarden LA, van Winkelhoff AJ. Activation of toll-like receptors 2 and 4 by gram-negative periodontal bacteria. *Oral Microbiol Immunol* 2007; **22**: 145–151.
 83. Kim S, Elkon KB, Ma X. Transcriptional suppression of interleukin-12 gene expression following phagocytosis of apoptotic cells. *Immunity* 2004; **21**: 643–653.
 84. Kim HM, Park BS, Kim JI, Kim SE, Lee J, Oh SC, Enkhbayar P, Matsushima N, Lee H, Yoo OJ, Lee JO. Crystal structure of the TLR4-MD-2 complex with bound endotoxin antagonist Eritoran. *Cell* 2007; **130**: 906–917.
 85. Kimizuka R, Kato T, Ishihara K, Okuda K. Mixed infections with *Porphyromonas gingivalis* and *Treponema denticola* cause excessive inflammatory responses in a mouse pneumonia model compared with mono-infections. *Microbes Infect* 2003; **5**: 1357–1362.
 86. Kimura Y, Miwa T, Zhou L, Song W-C. Activator-specific requirement of properdin in the initiation and amplification of the alternative pathway complement. *Blood* 2008; **111**: 732–740.
 87. Kinane DF, Demuth DR, Gorr SU, Hajishengallis GN, Martin MH. Human variability in innate immunity. *Periodontol 2000* 2007; **45**: 14–34.
 88. Kinane DF, Hajishengallis G. Polymicrobial infections, biofilms, and beyond. *J Clin Periodontol* 2009; **36**: 404–405.
 89. Kinane DF, Peterson M, Stathopoulou PG. Environmental and other modifying factors of the periodontal diseases. *Periodontol 2000* 2006; **40**: 107–119.
 90. Kline KA, Falker S, Dahlberg S, Normark S, Henriques-Normark B. Bacterial adhesins in host-microbe interactions. *Cell Host Microbe* 2009; **5**: 580–592.
 91. Klos A, Tenner AJ, Johswich KO, Ager RR, Reis ES, Kohl J. The role of the anaphylatoxins in health and disease. *Mol Immunol* 2009; **46**: 2753–2766.
 92. Kolenbrander PE, Parrish KD, Andersen RN, Greenberg EP. Intergeneric coaggregation of oral *Treponema* spp. with *Fusobacterium* spp. and intrageneric coaggregation among *Fusobacterium* spp. *Infect Immun* 1995; **63**: 4584–4588.
 93. Kumpf O, Schumann RR. Genetic influence on bloodstream infections and sepsis. *Int J Antimicrob Agents* 2008; **32** (Suppl. 1): S44–S50.

94. Kusumoto Y, Hirano H, Saitoh K, Yamada S, Takedachi M, Nozaki T, Ozawa Y, Nakahira Y, Saho T, Ogo H, Shimabukuro Y, Okada H, Murakami S. Human gingival epithelial cells produce chemotactic factors interleukin-8 and monocyte chemoattractant protein-1 after stimulation with *Porphyromonas gingivalis* via toll-like receptor 2. *J Periodontol* 2004; **75**: 370–379.
95. Lambris JD. *Current topics in complement II*. New York, NY: Springer, 2008.
96. Lambris JD, Ricklin D, Geisbrecht BV. Complement evasion by human pathogens. *Nat Rev Microbiol* 2008; **6**: 132–142.
97. Lamont RJ, Jenkinson HF. Life below the gum line: pathogenic mechanisms of *Porphyromonas gingivalis*. *Microbiol Mol Biol Rev* 1998; **62**: 1244–1263.
98. Lappegård KT, Christiansen D, Pharo A, Thorgersen BE, Hellerud BC, Lindstad J, Nielsen EW, Bergseth G, Fadnes D, Abrahamsen TG, Høyby A, Schejbel L, Garred P, Lambris JDP, Harboe M, Mollnes TE. Human genetic deficiencies reveal the roles of complement in the inflammatory network: lessons from nature. *Proc Natl Acad Sci USA* 2009; in press.
99. Lemaitre B, Nicolas E, Michaut L, Reichart JM, Hoffmann JA. The dorsoventral regulatory gene cassette *spatzle/Toll/cactus* controls the potent antifungal response in *Drosophila* adults. *Cell* 1996; **86**: 973–983.
100. Liew FY, Xu D, Brint EK, O'Neill LAJ. Negative regulation of Toll-like receptor-mediated immune responses. *Nat Rev Immunol* 2005; **5**: 446–458.
101. Lubberts E. IL-17/Th17 targeting: on the road to prevent chronic destructive arthritis? *Cytokine* 2008; **41**: 84–91.
102. Magalhaes MA, Sun CX, Glogauer M, Ellen RP. The major outer sheath protein of *Treponema denticola* selectively inhibits Rac1 activation in murine neutrophils. *Cell Microbiol* 2008; **10**: 344–354.
103. Markiewski MM, Lambris JD. The role of complement in inflammatory diseases from behind the scenes into the spotlight. *Am J Pathol* 2007; **171**: 715–727.
104. McDowell JV, Huang B, Fenno JC, Marconi RT. Analysis of a unique interaction between the complement regulatory protein factor H and the periodontal pathogen *Treponema denticola*. *Infect Immun* 2009; **77**: 1417–1425.
105. Medzhitov R. Recognition of microorganisms and activation of the immune response. *Nature* 2007; **449**: 819–826.
106. Medzhitov R, Preston-Hurlburt P, Janeway CA. A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature* 1997; **388**: 394–397.
107. Mevorach D, Mascarenhas JO, Gershov D, Elkon KB. Complement-dependent clearance of apoptotic cells by human macrophages. *J Exp Med* 1998; **188**: 2313–2320.
108. Millin DJ, Smith MH. Nature and composition of dental plaque. *Nature* 1961; **189**: 664–665.
109. Montminy SW, Khan N, McGrath S, Walkowicz MJ, Sharp F, Conlon JE, Fukase K, Kusumoto S, Sweet C, Miyake K, Akira S, Cotter RJ, Goguen JD, Lien E. Virulence factors of *Yersinia pestis* are overcome by a strong lipopolysaccharide response. *Nat Immunol* 2006; **7**: 1066–1073.
110. Morgan BP. Regulation of the complement membrane attack pathway. *Crit Rev Immunol* 1999; **19**: 173–198.
111. Mori Y, Yoshimura A, Ukai T, Lien E, Espevik T, Hara Y. Immunohistochemical localization of Toll-like receptors 2 and 4 in gingival tissue from patients with periodontitis. *Oral Microbiol Immunol* 2003; **18**: 54–58.
112. Muthukuru M, Jotwani R, Cutler CW. Oral mucosal endotoxin tolerance induction in chronic periodontitis. *Infect Immun* 2005; **73**: 687–694.
113. Mydel P, Takahashi Y, Yumoto H, Sztukowska M, Kubica M, Gibson FC 3rd, Kurtz DM Jr, Travis J, Collins LV, Nguyen KA, Genco CA, Potempa J. Roles of the host oxidative immune response and bacterial antioxidant rubrerythrin during *Porphyromonas gingivalis* infection. *PLoS Pathog* 2006; **2**: e76 (712–725).
114. Namavar F, Verweij AM, Bal M, van Steenberghe TJ, de Graaff J, MacLaren DM. Effect of anaerobic bacteria on killing of *Proteus mirabilis* by human polymorphonuclear leukocytes. *Infect Immun* 1983; **40**: 930–935.
115. Natarajan M, Lin KM, Hsueh RC, Sternweis PC, Ranganathan R. A global analysis of cross-talk in a mammalian cellular signalling network. *Nat Cell Biol* 2006; **8**: 571–580.
116. Nordahl EA, Rydengard V, Nyberg P, Nitsche DP, Morgelin M, Malmsten M, Björck L, Schmidtchen A. Activation of the complement system generates antibacterial peptides. *Proc Natl Acad Sci USA* 2004; **101**: 16879–16884.
117. Nussbaum G, Ben-Adi S, Genzler T, Sela M, Rosen G. TLR2 and TLR4 involvement in the innate immune response to *Treponema denticola* and its outer sheath components. *Infect Immun* 2009; **77**: 3939–3947.
118. O'Neill LA. When signaling pathways collide: positive and negative regulation of toll-like receptor signal transduction. *Immunity* 2008; **29**: 12–20.
119. O'Neill LA, Bowie AG. The family of five: TIR-domain-containing adaptors in Toll-like receptor signalling. *Nat Rev Immunol* 2007; **7**: 353–364.
120. Okuda K, Yanagi K, Takazoe I. Complement activation by *Propionibacterium acnes* and *Bacteroides melaninogenicus*. *Arch Oral Biol* 1978; **23**: 911–915.
121. Onishi S, Honma K, Liang S, Stathopoulou P, Kinane D, Hajishengallis G, Sharma A. Toll-like receptor 2-mediated interleukin-8 expression in gingival epithelial cells by the *Tannerella forsythia* leucine-rich repeat protein BspA. *Infect Immun* 2008; **76**: 198–205.
122. Pai RK, Convery M, Hamilton TA, Boom WH, Harding CV. Inhibition of IFN-gamma-induced class II transactivator expression by a 19-kDa lipoprotein from *Mycobacterium tuberculosis*: a potential mechanism for immune evasion. *J Immunol* 2003; **171**: 175–184.
123. Patters MR, Niekrash CE, Lang NP. Assessment of complement cleavage in gingival fluid during experimental gingivitis in man. *J Clin Periodontol* 1989; **16**: 33–37.
124. Pecora ND, Gehring AJ, Canaday DH, Boom WH, Harding CV. *Mycobacterium tuberculosis* LprA is a lipoprotein agonist of TLR2 that regulates innate immunity and APC function. *J Immunol* 2006; **177**: 422–429.
125. Pihlstrom BL, Michalowicz BS, Johnson NW. Periodontal diseases. *Lancet* 2005; **366**: 1809–1820.
126. Polak D, Wilensky A, Shapira L, Halabi A, Goldstein D, Weiss EI, Houry-Haddad Y. Mouse model of experimental periodontitis induced by *P. gingivalis* / *F. nucleatum* infection: bone loss and host response. *J Clin Periodontol* 2009; **36**: 406–410.
127. Popadiak K, Potempa J, Riesbeck K, Blom AM. Biphasic effect of gingipains from *Porphyromonas gingivalis* on the

- human complement system. *J Immunol* 2007; **178**: 7242–7250.
128. Potempa J, Golonka E, Filipek R, Shaw LN. Fighting an enemy within: cytoplasmic inhibitors of bacterial cysteine proteases. *Mol Microbiol* 2005; **57**: 605–610.
129. Potempa J, Pike RN. Corruption of innate immunity by bacterial proteases. *J Innate Immun* 2009; **1**: 70–87.
130. Potempa M, Potempa J, Kantyka T, Nguyen KA, Wawrzonek K, Manandhar SP, Popadiak K, Riesbeck K, Eick S, Blom AM. Interpain A, a cysteine proteinase from *Prevotella intermedia*, inhibits complement by degrading complement factor C3. *PLoS Pathog* 2009; **5**: e1000316.
131. Potempa M, Potempa J, Okroj M, Popadiak K, Eick S, Nguyen KA, Riesbeck K, Blom AM. Binding of complement inhibitor C4b-binding protein contributes to serum resistance of *Porphyromonas gingivalis*. *J Immunol* 2008; **181**: 5537–5544.
132. Pulendran B, Kumar P, Cutler CW, Mohamadzadeh M, Van Dyke T, Banchereau J. Lipopolysaccharides from distinct pathogens induce different classes of immune responses *in vivo*. *J Immunol* 2001; **167**: 5067–5076.
133. Re F, Strominger JL. Toll-like receptor 2 (TLR2) and TLR4 differentially activate human dendritic cells. *J Biol Chem* 2001; **276**: 37692–37699.
134. Ren L, Leung WK, Darveau RP, Jin L. The expression profile of lipopolysaccharide-binding protein, membrane-bound CD14, and toll-like receptors 2 and 4 in chronic periodontitis. *J Periodontol* 2005; **76**: 1950–1959.
135. Ricklin D, Lambris JD. Complement-targeted therapeutics. *Nat Biotechnol* 2007; **25**: 1265–1275.
136. Riedemann NC, Guo RF, Bernacki KD, Reuben JS, Laudes IJ, Neff TA, Gao H, Speyer C, Sarma VJ, Zetoune FS, Ward PA. Regulation by C5a of neutrophil activation during sepsis. *Immunity* 2003; **19**: 193–202.
137. Rittirsch D, Flierl MA, Ward PA. Harmful molecular mechanisms in sepsis. *Nat Rev Immunol* 2008; **8**: 776–787.
138. Rosen G, Genzler T, Sela MN. Coaggregation of *Treponema denticola* with *Porphyromonas gingivalis* and *Fusobacterium nucleatum* is mediated by the major outer sheath protein of *Treponema denticola*. *FEMS Microbiol Lett* 2008; **289**: 59–66.
139. Rosenberger CM, Finlay BB. Phagocyte sabotage: disruption of macrophage signalling by bacterial pathogens. *Nat Rev Mol Cell Biol* 2003; **4**: 385–396.
140. Roy CR, Mocarski ES. Pathogen subversion of cell-intrinsic innate immunity. *Nat Immunol* 2007; **8**: 1179–1187.
141. Ruby J, Rehani K, Martin M. *Treponema denticola* activates mitogen-activated protein kinase signal pathways through Toll-like receptor 2. *Infect Immun* 2007; **75**: 5763–5768.
142. Sabroe I, Dower SK, Whyte MK. The role of Toll-like receptors in the regulation of neutrophil migration, activation, and apoptosis. *Clin Infect Dis* 2005; **41** (Suppl. 7): S421–S426.
143. la Sala A, Gadina M, Kelsall BL. G(i)-protein-dependent inhibition of IL-12 production is mediated by activation of the phosphatidylinositol 3-kinase-protein 3 kinase B/Akt pathway and JNK. *J Immunol* 2005; **175**: 2994–2999.
144. Sato K. Th17 cells and rheumatoid arthritis—from the standpoint of osteoclast differentiation. *Allergol Int* 2008; **57**: 109–114.
145. Schenkein HA. Complement factor D-like activity of *Porphyromonas gingivalis* W83. *Oral Microbiol Immunol* 1991; **6**: 216–220.
146. Schenkein HA, Berry CR. Activation of complement by *Treponema denticola*. *J Dent Res* 1991; **70**: 107–110.
147. Schenkein HA, Genco RJ. Gingival fluid and serum in periodontal diseases. I. Quantitative study of immunoglobulins, complement components, and other plasma proteins. *J Periodontol* 1977; **48**: 772–777.
148. Schenkein HA, Genco RJ. Gingival fluid and serum in periodontal diseases. II. Evidence for cleavage of complement components C3, C3 proactivator (factor B) and C4 in gingival fluid. *J Periodontol* 1977; **48**: 778–784.
149. Sela MN. Role of *Treponema denticola* in periodontal diseases. *Crit Rev Oral Biol Med* 2001; **12**: 399–413.
150. Sela MN, Weinberg A, Borinsky R, Holt SC, Dishon T. Inhibition of superoxide production in human polymorphonuclear leukocytes by oral treponemal factors. *Infect Immun* 1988; **56**: 589–594.
151. Shibata Y, Fujimura S, Nakamura T. Purification and partial characterization of an elastolytic serine protease of *Prevotella intermedia*. *Appl Environ Microbiol* 1993; **59**: 2107–2111.
152. Shin H, Mally M, Kuhn M, Paroz C, Cornelis GR. Escape from immune surveillance by *Capnocytophaga canimorsus*. *J Infect Dis* 2007; **195**: 375–386.
153. Silverstein AM. *Paul Ehrlich's receptor immunology*. San Diego: Academic Press, 2002.
154. Sing A, Reithmeier-Rost D, Granfors K, Hill J, Roggenkamp A, Heesemann J. A hypervariable N-terminal region of *Yersinia LcrV* determines Toll-like receptor 2-mediated IL-10 induction and mouse virulence. *Proc Natl Acad Sci USA* 2005; **102**: 16049–16054.
155. Slaney JM, Curtis MA. Mechanisms of evasion of complement by *Porphyromonas gingivalis*. *Front Biosci* 2008; **13**: 188–196.
156. Slaney JM, Gallagher A, Aduse-Opoku J, Pell K, Curtis MA. Mechanisms of resistance of *Porphyromonas gingivalis* to killing by serum complement. *Infect Immun* 2006; **74**: 5352–5361.
157. Socransky SS, Haffajee AD. Periodontal microbial ecology. *Periodontol* 2000 2005; **38**: 135–187.
158. Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent RL Jr. Microbial complexes in subgingival plaque. *J Clin Periodontol* 1998; **25**: 134–144.
159. Sugawara S, Yang S, Iki K, Hatakeyama J, Tamai R, Takeuchi O, Akashi S, Espevik T, Akira S, Takada H. Monocytic cell activation by Nonendotoxic glycoprotein from *Prevotella intermedia* ATCC 25611 is mediated by toll-like receptor 2. *Infect Immun* 2001; **69**: 4951–4957.
160. Sunde PT, Olsen I, Gobel UB, Theegarten D, Winter S, Debelian GJ, Tronstad L, Moter A. Fluorescence *in situ* hybridization (FISH) for direct visualization of bacteria in periapical lesions of asymptomatic root-filled teeth. *Microbiology* 2003; **149**: 1095–1102.
161. Sundqvist G, Johansson E. Bactericidal effect of pooled human serum on *Bacteroides melaninogenicus*, *Bacteroides asaccharolyticus* and *Actinobacillus actinomycetemcomitans*. *Scand J Dent Res* 1982; **90**: 29–36.
162. Tabeta K, Yamazaki K, Akashi S, Miyake K, Kumada H, Umemoto T, Yoshie H. Toll-like receptors confer

- responsiveness to lipopolysaccharide from *Porphyromonas gingivalis* in human gingival fibroblasts. *Infect Immun* 2000; **68**: 3731–3735.
163. Tamai R, Deng X, Kiyoura Y. *Porphyromonas gingivalis* with either *Tannerella forsythia* or *Treponema denticola* induces synergistic IL-6 production by murine macrophage-like J774.1 cells. *Anaerobe* 2009; **15**: 87–90.
164. Tanner AC, Izard J. *Tannerella forsythia*, a periodontal pathogen entering the genomic era. *Periodontol* 2000 2006; **42**: 88–113.
165. Tanner A, Listgarten M, Ebersole J, Strzempko M. *Bacteroides forsythus* sp. nov., a slow-growing, fusiform *Bacteroides* sp. from the human oral cavity. *Int J Syst Bacteriol* 1986; **35**: 213–221.
166. Teng YT. Protective and destructive immunity in the periodontium: Part 1-Innate and humoral immunity and the periodontium. *J Dent Res* 2006; **85**: 198–208.
167. Toft RW, Peterson PK, Schmeling D, Bracke J, Kim Y, Quie PG. Opsonization of four *Bacteroides* species: role of the classical complement pathway and immunoglobulin. *Infect Immun* 1980; **27**: 784–792.
168. Trinchieri G. Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nat Rev Immunol* 2003; **3**: 133–146.
169. Ukai T, Yumoto H, Gibson FC 3rd, Genco CA. Macrophage-elicited osteoclastogenesis in response to bacterial stimulation requires Toll-like receptor 2-dependent tumor necrosis factor- α production. *Infect Immun* 2008; **76**: 812–819.
170. Wang PL, Ohura K, Fujii T, Oido-Mori M, Kowashi Y, Kikuchi M, Suetsugu Y, Tanaka J. DNA microarray analysis of human gingival fibroblasts from healthy and inflammatory gingival tissues. *Biochem Biophys Res Commun* 2003; **305**: 970–973.
171. Wang M, Shakhathreh M-AK, James D, Liang S, Nishiyama S-i, Yoshimura F, Demuth DR, Hajishengallis G. Fimbrial proteins of *Porphyromonas gingivalis* mediate *in vivo* virulence and exploit TLR2 and complement receptor 3 to persist in macrophages. *J Immunol* 2007; **179**: 2349–2358.
172. Ward PA. The dark side of C5a in sepsis. *Nat Rev Immunol* 2004; **4**: 133–142.
173. Ward PA. Functions of C5a receptors. *J Mol Med* 2009; **87**: 375–378.
174. Watanabe I, Ichiki M, Shiratsuchi A, Nakanishi Y. TLR2-mediated survival of *Staphylococcus aureus* in macrophages: a novel bacterial strategy against host innate immunity. *J Immunol* 2007; **178**: 4917–4925.
175. Weiss-Haljiti C, Pasquali C, Ji H, Gillieron C, Chabert C, Curchod ML, Hirsch E, Ridley AJ, Hooft van Huijsduijnen R, Camps M, Rommel C. Involvement of phosphoinositide 3-kinase gamma, Rac, and PAK signaling in chemokine-induced macrophage migration. *J Biol Chem* 2004; **279**: 43273–43284.
176. Wingrove JA, DiScipio RG, Chen Z, Potempa J, Travis J, Hugli TE. Activation of complement components C3 and C5 by a cysteine proteinase (gingipain-1) from *Porphyromonas (Bacteroides) gingivalis*. *J Biol Chem* 1992; **267**: 18902–18907.
177. Wright SD, Silverstein SC. Receptors for C3b and C3bi promote phagocytosis but not the release of toxic oxygen from human phagocytes. *J Exp Med* 1983; **158**: 2016–2023.
178. Ximenez-Fyvie LA, Haffajee AD, Socransky SS. Microbial composition of supra- and subgingival plaque in subjects with adult periodontitis. *J Clin Periodontol* 2000; **27**: 722–732.
179. Yamazaki T, Miyamoto M, Yamada S, Okuda K, Ishihara K. Surface protease of *Treponema denticola* hydrolyzes C3 and influences function of polymorphonuclear leukocytes. *Microbes Infect* 2006; **8**: 1758–1763.
180. Yanagisawa M, Kuriyama T, Williams DW, Nakagawa K, Karasawa T. Proteinase activity of *Prevotella* species associated with oral purulent infection. *Curr Microbiol* 2006; **52**: 375–378.
181. Yilmaz O. The chronicles of *Porphyromonas gingivalis*: the microbium, the human oral epithelium and their interplay. *Microbiology* 2008; **154**: 2897–2903.
182. Yoshimura A, Kaneko T, Kato Y, Golenbock DT, Hara Y. Lipopolysaccharides from periodontopathic bacteria *Porphyromonas gingivalis* and *Capnocytophaga ochracea* are antagonists for human toll-like receptor 4. *Infect Immun* 2002; **70**: 218–225.
183. Zhang X, Kimura Y, Fang C, Zhou L, Sfyroera G, Lambris JD, Wetsel RA, Miwa T, Song WC. Regulation of Toll-like receptor-mediated inflammatory response by complement *in vivo*. *Blood* 2007; **110**: 228–236.